

Hydrogen gas protects against serum and glucose deprivation-induced myocardial injury in H9c2 cells through activation of the NF-E2-related factor 2/heme oxygenase 1 signaling pathway

QIANG XIE¹, XUE-XIANG LI², PENG ZHANG¹, JIN-CAO LI¹, YING CHENG¹,
YAN-LING FENG¹, BING-SHENG HUANG¹, YU-FENG ZHUO¹ and GUO-HUA XU¹

¹Department of Cardiology, Hexian Memorial Hospital, Panyu, Guangzhou, Guangdong 511400;

²Department of Cardiology, Laizhou People's Hospital, Laizhou, Shandong 261400, P.R. China

Received September 16, 2013; Accepted April 25, 2014

DOI: 10.3892/mmr.2014.2283

Abstract. Ischemia or hypoxia-induced myocardial injury is closely associated with oxidative stress. Scavenging free radicals and/or enhancing endogenous antioxidative defense systems may be beneficial for the impediment of myocardial ischemic injury. Hydrogen (H₂) gas, as a water- and lipid-soluble small molecule, is not only able to selectively eliminate hydroxyl (\cdot OH) free radicals, but also to enhance endogenous antioxidative defense systems in rat lungs and arabidopsis plants. However, thus far, it has remained elusive whether H₂ gas protects cardiomyocytes through enhancement of endogenous antioxidative defense systems. In the present study, the cardioprotective effect of H₂ gas against ischemic or hypoxic injury was investigated, along with the underlying molecular mechanisms. H9c2 cardiomyoblasts (H9c2 cells) were treated *in vitro* with a chemical hypoxia inducer, cobalt chloride (CoCl₂), to imitate hypoxia, or by serum and glucose deprivation (SGD) to imitate ischemia. Cell viability and intracellular \cdot OH free radicals were assessed. The role of an endogenous antioxidative defense system, the NF-E2-related factor 2 (Nrf2)/heme oxygenase 1 (HO-1) signaling pathway, was evaluated. The findings revealed that treatment with CoCl₂ or SGD markedly reduced cell viability in H9c2 cells. H₂ gas-rich medium protected against cell injury induced by SGD, but not that induced by CoCl₂. When the cells were exposed to SGD, levels of intracellular \cdot OH free radicals were markedly increased; this was mitigated by H₂ gas-rich medium. Exposure of the cells to SGD also resulted in signifi-

cant increases in HO-1 expression and nuclear Nrf2 levels, and the HO-1 inhibitor ZnPP IX and the Nrf2 inhibitor brusatol aggravated SGD-induced cellular injury. H₂ gas-rich medium enhanced SGD-induced upregulation of HO-1 and Nrf2, and the HO-1 or Nrf2 inhibition partially suppressed H₂ gas-induced cardioprotection. Furthermore, following genetic silencing of Nrf2 by RNA interference, the effects of H₂ gas on the induction of HO-1 and cardioprotection were markedly reduced. In conclusion, H₂ gas protected cardiomyocytes from ischemia-induced myocardial injury through elimination of \cdot OH free radicals and also through activation of the Nrf2/HO-1 signaling pathway.

Introduction

Myocardial ischemic injury is a common pathological process in patients suffering from cardiac conditions, including atherosclerotic coronary artery disease, acute myocardial infarction and cardiac transplantation (1). However, the treatment costs for these diseases are high; therefore, it is important to investigate low-cost therapies that impede ischemia-induced myocardial injury.

The mechanisms underlying ischemia-induced cell damage are complicated and remain elusive. Increasing evidence suggested that oxidative stress is important in myocardial ischemic injury (2). Oxidative stress is characterized by marked increases in the production of reactive oxygen species (ROS), including the superoxide anion (O₂⁻) and the hydroxyl radical (\cdot OH), as well as non-radical molecules, including hydrogen peroxide (H₂O₂) and singlet oxygen (¹O₂) (3,4). Environmental stress factors, including ultraviolet rays, heat exposure, as well as ischemia and/or hypoxia, are major causes of oxidative stress, which may damage proteins, lipids and DNA, and eventually result in cellular death or the development of cancer (5,6). Under normal conditions, intracellular ROS levels are controlled by balancing ROS generation with ROS elimination. Once the balance is disrupted, for instance, through increased ROS generation and/or reduced elimination, ROS may aggregate and oxidative stress arises. Eliminating excessive ROS and enhancing endogenous antioxidation ability

Correspondence to: Professor Guo-Hua Xu, Department of Cardiology, Hexian Memorial Hospital, 2 Eastern Qinghe Road, Panyu, Guangzhou, Guangdong 511400, P.R. China
E-mail: ghxu2008@163.com

Key words: hydrogen gas, heme oxygenase 1, hydroxyl free radical, cardioprotection, NF-E2-related factor 2, serum and glucose deprivation

have been applied clinically (7), and in myocardial ischemic injury, oxidant scavengers, antioxidant extracts, vitamin E and vitamin C have all demonstrated to have a potential therapeutic value (8). However, water-soluble vitamin C has a low transmembrane diffusion ability and it is difficult to accumulate vitamin C up to an effective level to eliminate ROS (9). Conversely, the lipid-soluble vitamin E is difficult to dissolve in the cytoplasm in order to neutralize ROS (10). These shortcomings have limited the wide clinical use of the two vitamins. Therefore, a small antioxidant with water- and lipid-solubility is expected to have greater application value.

Hydrogen (H₂) gas is an inexpensive medical gas generated by electrolysis of water. Similar to other gaseous molecules, including nitric oxide (NO), carbon monoxide (CO) and hydrogen sulfide (H₂S), H₂ gas has been demonstrated to exhibit numerous important cytoprotective effects in the nervous, cardiovascular and digestive systems (11-14). Unlike vitamin C and vitamin E, H₂ gas dissolves in water and lipids. In addition, the simple molecular structure and small molecular weight render H₂ gas a good antioxidant candidate in cells. A number of studies have suggested that H₂ gas may selectively scavenge ·OH free radicals (15,16). Therefore, H₂ gas may have broad clinical applications in the future. However, in cardiomyocytes, it has not been reported whether H₂ gas protects against ischemia-induced injury *in vitro*, which was the focus of the present study.

Abundant evidence indicated that enhancement of endogenous antioxidation activity exerts cardioprotective effects. For instance, upregulation of superoxide dismutase (SOD) or heme oxygenase-1 (HO-1) protects cardiomyocytes against ischemia and/or reperfusion-induced damage (17-19). Notably, in addition to scavenging ·OH free radicals, H₂ gas protection has been associated with induction of HO-1 controlled by NF-E2-related factor 2 (Nrf2) in rat lung transplant-induced injury and paraquat-induced oxidative damage in plants (20,21). H₂ gas protection of cardiomyocytes may therefore be associated with scavenging ·OH free radicals and upregulation of the Nrf2/HO-1 signaling pathway.

H9c2 cardiomyoblasts (H9c2 cells), originating from rat heart ventricular tissue, have widely served as an *in vitro* model for cardiac muscle in virtue of their morphological features and biochemical properties (22). In the present study, H9c2 cells were subjected to serum and glucose deprivation (SGD) or exposed to hypoxia provoked by a chemical hypoxia-mimicking agent, cobalt chloride (CoCl₂), to establish an ischemia/hypoxia-induced myocardial injury model. H₂ gas-rich medium was applied to investigate the cytoprotection, antioxidation and activation of the Nrf2/HO-1 signaling pathway as well as the involvement of scavenging ·OH free radicals and upregulation of the Nrf2/HO-1 signaling pathway in the cardioprotection by H₂ gas.

Materials and methods

Materials. CoCl₂ and protoporphyrin IX zinc (II) (ZnPP IX) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Brusatol (BR), an inhibitor of Nrf2, was provided by BOC Sciences (Shirley, NY, USA). A cell counting kit-8 (CCK-8) was purchased from Dojindo Laboratories (Kyushu, Japan). Specific monoclonal primary antibodies against rat HO-1 and Nrf2 proteins were obtained from EPITOMICS of

Abcam Company (Burlingame, CA, USA). High glucose and glucose-free Dulbecco's modified Eagle's medium and fetal bovine serum (FBS) were supplied by Gibco-BRL (Grand Island, NY, USA). A bicinchoninic acid (BCA) protein assay kit was purchased from Kangchen Bio-tech Inc. (Shanghai, China). An enhanced chemiluminescence (ECL) kit was obtained from Applygen Technologies Inc. (Beijing, China). An enzyme-linked immunosorbent assay (ELISA) kit for detection of 8-hydroxy-2'-deoxyguanosine (8-OHdG) was provided by Abnova Corporation (Taipei, Taiwan).

Cell culture. H9c2 cells were supplied by the Cell Bank at the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). The cells were maintained in high glucose DMEM supplemented with 15% FBS at 37°C in an atmosphere of 5% CO₂ and 95% air. The cells were passaged approximately every two days after digestion with 0.2% trypsin.

Hypoxia or ischemia treatment. CoCl₂, a chemical hypoxia inducer, was co-incubated with H9c2 cells to induce hypoxia. An ischemia-induced myocardial injury model generated through SGD was prepared *in vitro* in the cell medium. The cell viability was used to indicate the extent of hypoxic or ischemic injury in the H9c2 cells.

Preparation of H₂ gas-rich medium. Pure H₂ gas (99.999% purity) was produced via electrolysis of water with a M177021 H₂ gas generator, supplied by Beijing Midwest Yuanda Technology Co., Ltd. (Beijing, China; 23). H₂ gas-rich medium was prepared freshly prior to saturating the medium with the generated H₂ gas for at least 30 min.

Determination of cell viability. Cell viability was analyzed by a CCK-8 assay following the manufacturer's instructions. H9c2 cells were plated in 96-well plates at a density of 5,000 cells/well. When the cells were grown to ~70% confluence, the indicated treatments were administered. At the end of the treatment, the CCK-8 solution (10 μl) at 1:10 dilution with FBS-free DMEM high glucose medium (100 μl) was added to each well followed by a further 3 h incubation at 37°C. Absorbance (A) was measured at 450 nm with a microplate reader produced by Molecular Devices, LLC (Sunnyvale, CA, USA). The mean A was used to calculate the percentage of cell viability according to the following equation: Percentage of viable cells = (A treatment group - A Blank group)/(A Control group - A Blank group) x 100%. Experiments were performed six times (24).

Western blot analysis of protein expression. H9c2 cells were plated in 60-mm diameter petri dishes. Following the indicated treatments, the cells were harvested and total proteins or nuclear proteins were extracted and quantified with the BCA kit and used to measure HO-1 or Nrf2 expression levels, respectively. Subsequent to denaturation by heating at 100°C for 5 min, equal quantities of proteins of the indicated groups were loaded and separated by 12% SDS-PAGE. The proteins in the gel were then transferred to a polyvinylidene fluoride membrane. Following blocking with 5% fat-free milk in Tris-buffered saline with Tween 20 (TBS-T), the membranes were incubated with rat monoclonal primary antibodies against HO-1 or Nrf2 overnight with gentle agitation at 4°C. β-actin and Lamin B served as




Figure 1. Effects of H₂ gas treatment on CoCl₂-induced injury in H9c2 cells. Following the indicated treatments, cell viability was measured by a cell counting kit-8 assay. (A) H9c2 cells were treated with increasing concentrations of CoCl₂ at 100-1,000 μM for 24 h. (B) H9c2 cells were treated with CoCl₂ at 400-800 μM for 24 h in the absence (H₂-Free) or presence (H₂-Rich) of H₂ gas application. Data are presented as mean ± standard deviation, n=6. *P<0.05, **P<0.01, compared with the control group.

loading controls. Subsequent to three washes with TBS-T, the membranes were incubated with anti-rabbit horseradish peroxidase-conjugated secondary antibodies at room temperature for 1 h. The membranes were washed again and developed with an ECL system. The membranes were then exposed to X-ray films (Kodak Company, Beijing, China). The integrated optical density of the protein bands was calculated by Image J 1.47 Software (National Institutes of Health, Bethesda, MD, USA).

Competitive ELISA for measurement of 8-OHdG. Intracellular ·OH free radicals cause oxidative damage to DNA to form 8-OHdG. Therefore, by measuring 8-OHdG levels in the cells, ·OH free radical content may be analyzed indirectly. H9c2 cells were plated in six-well plates and treated as indicated. At the end of the treatment, the cells were harvested and lysed. 8-OHdG levels in the lysate were determined according to the manufacturer's instructions (Abnova Corporation). The experiment was performed at least six times with similar outcomes.

Gene knockdown. Small interfering RNA (Si-RNA) against rat Nrf2 mRNA (GenBank accession no. AF037350; <https://www.ncbi.nlm.nih.gov/genbank/>) was synthesized by GenePharma Co., Ltd (Shanghai, China). The Si-RNA of Nrf2 (Si-Nrf2) and random non-coding RNA (Si-NC) were transfected into H9c2 cells using Lipofectamine 2000, according to the manufacturer's instructions (Invitrogen Life Technologies, Carlsbad, CA, USA). Si-Nrf2 and Si-NC (20 nmol/l) were incubated with the cells for 6 h followed by further incubation for 24 h in order to transfect the cells.

Statistical analysis. All data were analyzed using SPSS 13.0 software (SPSS, Inc., Chicago, IL, USA) and the results are expressed as the mean ± standard deviation. The significance of intergroup differences was evaluated by one-way analyses of variance. Differences were considered to be significant if the two-sided probability (P) was <0.05.

Results

H₂ gas-rich medium does not affect chemical hypoxia-induced myocardial injury. Hypoxemia can be imitated *in vitro* by

chemical hypoxia or SGD treatment. H₂ gas protection in chemical hypoxia-induced myocardial injury was firstly assessed by treatment of H9c2 cells with CoCl₂ in H₂ gas-rich medium. As shown in Fig. 1A, treatment of H9c2 cells with increasing concentrations of CoCl₂ significantly reduced cell viability at concentrations of 400-1,000 μM. In order to observe H₂ gas effectiveness, the cell culture medium was saturated with H₂ gas generated by electrolysis of water. No difference in cell viability between the cells cultivated in H₂ gas-rich medium and those cultivated in H₂ gas-free medium was identified under the conditions of CoCl₂ exposure or rest (Fig. 1B). This result indicated that H₂ gas treatment did not influence chemical hypoxia-induced myocardial injury.

H₂ gas-rich medium alleviates SGD-induced cell injury. SGD treatment acts as another *in vitro* hypoxemia model by withdrawing FBS and glucose to mimic acute infarction-induced myocardial injury *in vivo*. When the cells were exposed to SGD for different time periods followed by further culture for 30 h, the cell viability of H9c2 cells was significantly reduced (P<0.01; Fig. 2A). Following exposure to SGD for 6-18 h, the H9c2 cells were cultured in H₂ gas-rich medium for 30 h. The findings demonstrated that, compared with H₂ gas-free culture, H₂ gas-rich culture significantly increased the viability of the cells subjected to SGD for 6 or 12 h (P<0.05), but not 18 h. The results suggested that when the cell injury induced by SGD was not severe, H₂ gas was able to promote cell survival.

H₂ gas-rich medium prevents SGD-induced ·OH free radical generation. One mechanism underlying H₂ protection is ROS scavenging, particularly of ·OH free radicals. To investigate whether H₂ gas-mediated myocardial protection was associated with scavenging ·OH free radicals, the ·OH levels in H9c2 cells were measured using ELISA. As shown in Fig. 3, exposure of the cells to SGD significantly enhanced intracellular ·OH levels (P<0.01). However, this effectiveness was significantly inhibited by incubation in H₂ gas-rich medium (P<0.01). This indicated that the elimination of ·OH free radicals may be an important mechanism of myocardial protection by H₂ gas.

Adaptive HO-1 induction contributes to H₂ gas inhibition of SGD-induced injury in H9c2 cells. HO-1 is endogenously




Figure 2. Effects of H₂ gas on SGD-induced injury in H9c2 cells. Following the indicated treatments, cell viability was measured with a cell counting kit-8 assay. (A) H9c2 cells were exposed to SGD between 6 and 30 h, followed by a further 30 h in normal culture. (B) H9c2 cells were exposed to SGD for 6, 12 or 18 h, respectively, in the absence (H₂-Free) or presence (H₂-Rich) of H₂ gas treatment. Data are presented as the mean \pm standard deviation, n=6. *P<0.05, **P<0.01, compared with the control group. §P<0.05, compared with the SGD group. SGD, serum and glucose deprivation.




Figure 3. Effects of H₂ gas on SGD-induced hydroxyl free radical generation in H9c2 cells. The cells were exposed to SGD for 6 h in the absence or presence of H₂ gas application. Intracellular \cdot OH free radical content was indirectly measured by ELISA of 8-OHdG levels. Data are presented as the mean \pm standard deviation, n=6. *P<0.01, compared with the control group. §P<0.01, compared with the SGD group. SGD, serum and glucose deprivation; 8-OHdG, 8-hydroxy-2'-deoxyguanosine.

produced, functioning as an antioxidation enzyme. In order to examine the role of HO-1 in SGD-induced myocardial injury, experiments to detect HO-1 levels were performed. The data in Fig. 4 show that exposure of H9c2 cells to SGD significantly upregulated HO-1 levels when compared with the cells under normal conditions (P<0.01). When the HO-1 inhibitor ZnPP IX was administered, SGD-induced cellular injury was found to be significantly aggravated (P<0.05; Fig. 5). These results suggested that HO-1 was beneficial in SGD-induced injury in H9c2 cells.

Upregulation of HO-1 is involved in H₂ gas-induced myocardial protection. H₂ gas-induced protection is not only associated with simple elimination of \cdot OH free radicals, but also with induction of endogenous genes, for instance, HO-1 (20,21). To clarify whether HO-1 is involved in H₂ gas-induced myocardial protection, experiments were conducted to observe the effect of H₂ gas on SGD-triggered HO-1 upregulation and the effect of HO-1 inhibition on H₂ gas-induced protection of H9c2 cells. The data in Fig. 4 reveal that exogenously applied H₂ gas significantly enhanced the upregulation elicited by SGD in H9c2 cells (P<0.05). Notably, inhibition of HO-1 with 10 μ M




Figure 4. Levels of HO-1 under different conditions. H9c2 cells were exposed to SGD for 6 h followed by a further 30 h culture in H₂ gas-free/-rich medium. At the end of the treatments, HO-1 expression in the cells was detected by (A) Western blot assay and (B) quantitative analysis with ImageJ 1.47 software. Data are presented as the mean \pm standard deviation, n=6. *P<0.01 compared with the control group. §P<0.05 compared with the SGD group. SGD, serum and glucose deprivation; HO-1, heme oxygenase 1.

ZnPP IX significantly reduced the H₂ gas-induced increase in cellular viability following SGD treatment (P<0.05; Fig. 6). The results suggested that HO-1 at least partially mediated the protection from ischemia provided by H₂ gas in H9c2 cells.

H₂ gas facilitates nuclear Nrf2 expression induced by SGD in H9c2 cells. Nrf2 is a transcription factor responsible for HO-1 expression. To address the role of Nrf2 in H₂ gas-induced myocardial protection, the effect of H₂ gas-rich medium on the changes of Nrf2 levels induced by SGD was investigated. As shown in Fig. 7, the SGD challenge resulted in a significant increase in nuclear Nrf2 expression levels, indicating its activation under SGD conditions in H9c2 cells (P<0.05). Notably, H₂ gas application following SGD exposure induced a significant increase in nuclear Nrf2 expression levels (P<0.05).




Figure 5. Effects of ZnPP IX on SGD-induced injury in H9c2 cells. The cells were exposed to SGD for 6 h followed by further culture with or without an additional 10 μ M ZnPP IX treatment for 30 h. Following treatment, the cell viability was measured by the cell counting kit-8 assay. Data are presented as the mean \pm standard deviation, n=6. *P<0.01 compared with the control group. [§]P<0.05 compared with the SGD group. SGD, serum and glucose deprivation; ZnPP IX, protoporphyrin IX zinc (II).




Figure 6. Effect of the indicated treatments on myocardial viability. H9c2 cells were exposed to SGD for 6 h followed by a further culture combined with H₂ gas with or without 10 μ M ZnPP IX for 30 h. Following treatment, the cell viability was measured by the cell counting kit-8 assay. Data are presented as the mean \pm standard deviation, n=6. *P<0.01 compared with the control group. [§]P<0.05 compared with the SGD group. [#]P<0.05 compared with SGD combined with the H₂ gas group. SGD, serum and glucose deprivation; ZnPP IX, protoporphyrin IX zinc (II).

Inhibition of Nrf2 with 10 μ M BR significantly reduced myocardial protection by H₂ gas (P<0.05, Fig. 8).

In addition, genetic silencing of Nrf2 by RNAi (Si-Nrf2) also significantly inhibited H₂ gas-elicited HO-1 induction (P<0.05, Fig. 9) and myocardial protective action (P<0.05, Fig. 8). These results indicated that the H₂ gas protection from SGD-induced injury in H9c2 cells was at least in part mediated through the Nrf2/HO-1 signaling pathway.

Discussion

The results of the present study suggested that H₂ gas exhibited myocardial protection against ischemia-induced injury in H9c2 cells *in vitro* through elimination of \cdot OH free radicals and activation of the Nrf2/HO-1 signaling pathway. These findings provide further evidence of H₂ gas protection and deepen the understanding of the molecular mechanisms involved.




Figure 7. Effects of SGD and H₂ gas on expression of nuclear Nrf2 in H9c2 cells. The cells were exposed to SGD for 6 h followed by a further 30 h culture in H₂ gas-free/-rich medium. Following treatment, nuclear proteins were extracted and Nrf2 expression levels were detected by (A) western blot assay and (B) quantitative analysis with ImageJ 1.47 software. Data are presented as the mean \pm standard deviation, n=6. *P<0.05 compared with the control group. [§]P<0.05 compared with the SGD group. SGD, serum and glucose deprivation; Nrf2, NF-E2-related factor 2.




Figure 8. Effect of the different treatments on myocardial viability. Normal H9c2 cells were exposed to SGD for 6 h followed by a further 30 h culture combined with H₂ gas with or without 10 μ M BR. Nrf2-deficient H9c2 cells caused by RNA interference assay were exposed to SGD for 6 h followed by a further 30 h culture combined with H₂ gas. Following treatment, the cell viability was measured by a cell counting kit-8 assay. Data are presented as the mean \pm standard deviation, n=6. *P<0.01 compared with the control group. [§]P<0.05 compared with the SGD group. [#]P<0.05 compared with SGD combined with the H₂ gas group. SGD, serum and glucose deprivation; NC, negative control; BR, brusatol; Nrf2, NF-E2-related factor 2; Si, small interfering RNA.

H₂ gas is a gas with novel medical application, in addition to NO, CO and H₂S, whose cytoprotective effects have gradually gained attention (15). The cytoprotective effect of H₂ gas has been investigated in the nervous, cardiovascular and digestive systems (11-14). In the present study, myocardial protection by H₂ gas was investigated in two distinct models: Chemical hypoxia-induced injury and SGD-induced injury in H9c2 cells. Treatment with chemical hypoxia-mimicking




Figure 9. Effect of H₂ gas on the induction of HO-1 in Nrf2-deficient H9c2 cells. Following RNA interference with Si-Nrf2, which downregulated Nrf2 expression, the cells were exposed to SGD for 6 h followed by a further 30 h culture in the absence or presence of H₂ gas. Following treatment, HO-1 expression was observed by (A) western blot assay and (B) quantitative analysis with ImageJ 1.47 software. Data are presented as the mean \pm standard deviation, n=6. *P<0.01 compared with the control group. #P<0.05 compared with SGD combined with H₂ gas group. SGD, serum and glucose deprivation; Nrf2, NF-E2-related factor 2; Si, small interfering RNA; HO-1, heme oxygenase 1.

agent CoCl₂, at 400-1,000 μ M for 24 h, reduced cell viability in a concentration-dependent manner, although no effect of H₂ gas on CoCl₂-induced injury was observed. The effect of H₂ gas on SGD-induced injury was then analyzed. Notably, H₂ gas exerted marked myocardial protection, since its application impeded SGD-induced injury in H9c2 cells. These findings were in accordance with a study by Sun *et al* (29) on cardiac ischemia/reperfusion injury in a rat model. However, the findings of the present study also indicated that although CoCl₂ (25,26) and SGD (27) are frequently used in hypoxia/ischemia *in vitro* models, they may possess markedly different underlying injury mechanisms. In addition, as H₂ gas is a water- and lipid-soluble and simple molecule, H₂ gas may exhibit a greater clinical antioxidative value than the water-soluble vitamin C and lipid-soluble vitamin E.

Oxidative stress is critical in myocardial ischemic damage through overproduction of ROS (2). ROS in mammals include O₂⁻, H₂O₂ and \cdot OH. O₂⁻, H₂O₂ ROS are eliminated by corresponding enzymes. For example, O₂⁻ may be catalyzed into O₂ and H₂O₂ via dismutation (28), and H₂O₂, one of the most powerful oxidizers, may be converted into H₂O by catalase or guaiacol peroxidase, but may also be converted into \cdot OH. Although \cdot OH free radicals are toxic to cells, enzymes responsible for \cdot OH elimination remain to be identified. Therefore, it is important for antioxidants to eliminate \cdot OH free radicals and/or inhibit the production of \cdot OH. In the present study, exposure of H9c2 cells to SGD was found to significantly increase intracellular \cdot OH free radical levels as identified through assessment of 8-OHdG. The \cdot OH levels after SGD stimulation were significantly reduced during cell cultivation in H₂ gas-rich medium, in concurrence with previous reports (15,16,29).

Increasing evidence suggested that the cytoprotective effect of H₂ gas is not only associated with simple elimination of \cdot OH free radicals, but also with numerous signaling molecules (20,21). A study on plants demonstrated that pretreatment with H₂ gas enhanced the salt tolerance of arabidopsis through zinc-finger transcription factor ZAT10/12 (30). Additionally, the HO-1 signaling pathway has been observed to be involved in H₂ gas protection against paraquat-induced oxidative injury (21). In animals, Cai *et al* (31) revealed that H₂ gas treatment alleviated tumor necrosis factor-alpha-induced rat osteoblast inflammatory injury via upregulation of SOD activity. Furthermore, evidence has indicated that HO-1 induction mediated H₂ gas mitigation of rat lung injury resulting from transplantation (20). In the present study, SGD exposure was found to markedly increase HO-1 expression. The mechanism of HO-1 in SGD-induced myocardial insult was further elucidated using a selective HO-1 inhibitor, ZnPP IX. Since the data indicated that the addition of ZnPP IX markedly aggravated the SGD-induced insult, HO-1 may exhibit a protective action against ischemia, a hypothesis which is supported by the results of a study by Hwa *et al* (32). Notably, exogenously applied H₂ gas was found to result in a further increase in HO-1 expression; application of ZnPP IX partially abolished this H₂ gas-triggered myocardial protection. Therefore, H₂ gas protection may be partially mediated by HO-1 induction. One study suggested that inhalation of H₂ gas combined with CO, a product of HO-1, enhanced its therapeutic efficacy in ischemia/reperfusion-induced myocardial injury (33). These findings, alongside those of the present study, support the hypothesis that the mechanisms underlying H₂ gas-induced myocardial protection are not limited to the elimination of the \cdot OH free radical and may also include upregulation of protective genes.

Nrf2 belongs to the NF-E2 superfamily of nuclear basic leucine zipper transcription factors. Under conditions of oxidative stress or pharmacological stimuli, Nrf2, as an adaptive response, regulates phase II gene expression of numerous enzymes that serve to detoxify pro-oxidative stressors (34). In the promoter region of certain genes, such as HO-1 and SOD, Nrf2 binds to the cis-acting regulatory element or enhancer sequence and induces gene expression (35). In the present study, SGD exposure markedly induced nuclear Nrf2 expression, which was further enhanced by H₂ gas administration. When the action of Nrf2 was inhibited by BR, H₂ gas-induced myocardial protection was significantly attenuated. Studies have indicated that BR may inhibit the Nrf2 signaling pathway (36,37). In addition, genetic silencing of Nrf2 may also impede H₂-induced HO-1 expression and increases in cell viability. These data suggested that the activation of Nrf2 is involved in H₂ gas protection and HO-1 induction. One study observed that another medical gas, H₂S, protected cardiomyocytes against ischemia-induced injury by activation of the Nrf2 signaling pathway (38). Nrf2-mediated myocardial protection has also been reported regarding a number of other compounds (39,40). For instance, in the nervous system, curcumin protected rat brains against focal ischemia via upregulation of the Nrf2/HO-1 signaling pathway (41). Therefore, the upregulation of the Nrf2/HO-1 signaling pathway is considered to be a general antioxidative mechanism of numerous drugs and compounds, and this signaling pathway may represent a novel molecular target of drug design.

In conclusion, the present study suggested that H₂ gas application not only directly scavenged ·OH free radicals but also enhanced the expression of proteins of the Nrf2/HO-1 signaling pathway in SGD-stimulated H9c2 cells. These findings provided basic information for the development of novel treatments of ischemic myocardial injury with H₂ gas.

Acknowledgements

This study was supported by Science & Technology Planning Project of Guangdong Province in China (no. 2012A030400033).

References

- Shibata R, Sato K, Pimentel DR, *et al*: Adiponectin protects against myocardial ischemia-reperfusion injury through AMPK- and COX-2-dependent mechanisms. *Nat Med* 11: 1096-1103, 2005.
- Yang XY, Zhao N, Liu YY, *et al*: Inhibition of NADPH oxidase mediates protective effect of cardioprotective pills against rat heart ischemia/reperfusion injury. *Evid Based Complement Alternat Med* 2013: 728020, 2013.
- Dickinson BC and Chang CJ: Chemistry and biology of reactive oxygen species in signaling or stress responses. *Nat Chem Biol* 7: 504-511, 2011.
- Devasagayam TP, Tilak JC, Bolor KK, Sane KS, Ghaskadbi SS and Lele RD: Free radicals and antioxidants in human health: current status and future prospects. *J Assoc Physicians India* 52: 794-804, 2004.
- Martinez-Outschoorn UE, Balliet RM, Rivadeneira DB, *et al*: Oxidative stress in cancer associated fibroblasts drives tumor-stroma co-evolution: A new paradigm for understanding tumor metabolism, the field effect and genomic instability in cancer cells. *Cell Cycle* 9: 3256-3276, 2010.
- Ryter SW, Kim HP, Hoetzel A, *et al*: Mechanisms of cell death in oxidative stress. *Antioxid Redox Signal* 9: 49-89, 2007.
- Jha P, Flather M, Lonn E, Farkouh M and Yusuf S: The antioxidant vitamins and cardiovascular disease. A critical review of epidemiologic and clinical trial data. *Ann Intern Med* 123: 860-872, 1995.
- Hamilton KL: Antioxidants and cardioprotection. *Med Sci Sports Exerc* 39: 1544-1553, 2007.
- Padayatty SJ, Katz A, Wang Y, *et al*: Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr* 22: 18-35, 2003.
- Ingold KU, Webb AC, Witter D, Burton GW, Metcalfe TA and Muller DP: Vitamin E remains the major lipid-soluble, chain-breaking antioxidant in human plasma even in individuals suffering severe vitamin E deficiency. *Arch Biochem Biophys* 259: 224-225, 1987.
- Wang C, Li J, Liu Q, *et al*: Hydrogen-rich saline reduces oxidative stress and inflammation by inhibit of JNK and NF-κB activation in a rat model of amyloid-beta-induced Alzheimer's disease. *Neurosci Lett* 491: 127-132, 2011.
- Zheng X, Mao Y, Cai J, *et al*: Hydrogen-rich saline protects against intestinal ischemia/reperfusion injury in rats. *Free Radic Res* 43: 478-484, 2009.
- Liu Q, Shen WF, Sun HY, *et al*: Hydrogen-rich saline protects against liver injury in rats with obstructive jaundice. *Liver Int* 30: 958-968, 2010.
- Yu P, Wang Z, Sun X, *et al*: Hydrogen-rich medium protects human skin fibroblasts from high glucose or mannitol induced oxidative damage. *Biochem Biophys Res Commun* 409: 350-355, 2011.
- Ohsawa I, Ishikawa M, Takahashi K, *et al*: Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals. *Nat Med* 13: 688-694, 2007.
- Fukuda K, Asoh S, Ishikawa M, Yamamoto Y, Ohsawa I and Ohta S: Inhalation of hydrogen gas suppresses hepatic injury caused by ischemia/reperfusion through reducing oxidative stress. *Biochem Biophys Res Commun* 361: 670-674, 2007.
- Ambrosio G, Becker LC, Hutchins GM, Weisman HF and Weisfeldt ML: Reduction in experimental infarct size by recombinant human superoxide dismutase: insights into the pathophysiology of reperfusion injury. *Circulation* 74: 1424-1433, 1986.
- Zhang X, Xiao Z, Yao J, Zhao G, Fa X and Niu J: Participation of protein kinase C in the activation of Nrf2 signaling by ischemic preconditioning in the isolated rabbit heart. *Mol Cell Biochem* 372: 169-179, 2013.
- Hui Y, Zhao Y, Ma N, *et al*: M3-mAChR stimulation exerts anti-apoptotic effect via activating the HIF-1α/HO-1/VEGF signaling pathway in H9c2 rat ventricular cells. *J Cardiovasc Pharmacol* 60: 474-482, 2012.
- Kawamura T, Wakabayashi N, Shigemura N, *et al*: Hydrogen gas reduces hyperoxic lung injury via the Nrf2 pathway in vivo. *Am J Physiol Lung Cell Mol Physiol* 304: L646-L656, 2013.
- Jin Q, Zhu K, Cui W, Xie Y, Han B and Shen W: Hydrogen gas acts as a novel bioactive molecule in enhancing plant tolerance to paraquat-induced oxidative stress via the modulation of heme oxygenase-1 signalling system. *Plant Cell Environ* 36: 956-969, 2013.
- Branco AF, Pereira SL, Moreira AC, Holy J, Sardão VA and Oliveira PJ: Isoproterenol cytotoxicity is dependent on the differentiation state of the cardiomyoblast H9c2 cell line. *Cardiovasc Toxicol* 11: 191-203, 2011.
- Huang G, Zhou J, Zhan W, *et al*: The neuroprotective effects of intraperitoneal injection of hydrogen in rabbits with cardiac arrest. *Resuscitation* 84: 690-695, 2013.
- Yang C, Yang Z, Zhang M, *et al*: Hydrogen sulfide protects against chemical hypoxia-induced cytotoxicity and inflammation in HaCaT cells through inhibition of ROS/NF-κB/COX-2 pathway. *PLoS One* 6: e21971, 2011.
- Chen SL, Yang CT, Yang ZL, *et al*: Hydrogen sulphide protects H9c2 cells against chemical hypoxia-induced injury. *Clin Exp Pharmacol Physiol* 37: 316-321, 2010.
- Yang Z, Yang C, Xiao L, *et al*: Novel insights into the role of HSP90 in cytoprotection of H₂S against chemical hypoxia-induced injury in H9c2 cardiac myocytes. *Int J Mol Med* 28: 397-403, 2011.
- Yao LL, Wang YG, Cai WJ, Yao T and Zhu YC: Survivin mediates the anti-apoptotic effect of delta-opioid receptor stimulation in cardiomyocytes. *J Cell Sci* 120: 895-907, 2007.
- Elchuri S, Oberley TD, Qi W, *et al*: CuZnSOD deficiency leads to persistent and widespread oxidative damage and hepatocarcinogenesis later in life. *Oncogene* 24: 367-380, 2005.
- Sun Q, Kang Z, Cai J, *et al*: Hydrogen-rich saline protects myocardium against ischemia/reperfusion injury in rats. *Exp Biol Med (Maywood)* 234: 1212-1219, 2009.
- Xie Y, Mao Y, Lai D, Zhang W and Shen W: H₂ enhances arbidopsis salt tolerance by manipulating ZAT10/12-mediated antioxidant defence and controlling sodium exclusion. *PLoS One* 7: e49800, 2012.
- Cai WW, Zhang MH, Yu YS and Cai JH: Treatment with hydrogen molecule alleviates TNFα-induced cell injury in osteoblast. *Mol Cell Biochem* 373: 1-9, 2013.
- Hwa JS, Jin YC, Lee YS, *et al*: 2-methoxycinnamaldehyde from *Cinnamomum cassia* reduces rat myocardial ischemia and reperfusion injury in vivo due to HO-1 induction. *J Ethnopharmacol* 139: 605-615, 2012.
- Nakao A, Kaczorowski DJ, Wang Y, *et al*: Amelioration of rat cardiac cold ischemia/reperfusion injury with inhaled hydrogen or carbon monoxide, or both. *J Heart Lung Transplant* 29: 544-553, 2010.
- Fisher CD, Augustine LM, Maher JM, *et al*: Induction of drug-metabolizing enzymes by garlic and allyl sulfide compounds via activation of constitutive androstane receptor and nuclear factor E2-related factor 2. *Drug Metab Dispos* 35: 995-1000, 2007.
- Kang KW, Lee SJ and Kim SG: Molecular mechanism of nrf2 activation by oxidative stress. *Antioxid Redox Signal* 7: 1664-1673, 2005.
- Bauer AK, Hill T III and Alexander CM: The involvement of NRF2 in lung cancer. *Oxid Med Cell Longev* 2013: 746432, 2013.
- Ren D, Villeneuve NF, Jiang T, *et al*: Brusatol enhances the efficacy of chemotherapy by inhibiting the Nrf2-mediated defense mechanism. *Proc Natl Acad Sci USA* 108: 1433-1438, 2011.
- Calvert JW, Jha S, Gundewar S, *et al*: Hydrogen sulfide mediates cardioprotection through Nrf2 signaling. *Circ Res* 105: 365-374, 2009.
- Deng C, Sun Z, Tong G, *et al*: α-Lipoic acid reduces infarct size and preserves cardiac function in rat myocardial ischemia/reperfusion injury through activation of PI3K/Akt/Nrf2 pathway. *PLoS One* 8: e58371, 2013.
- Cui G, Shan L, Hung M, *et al*: A novel Danshensu derivative confers cardioprotection via PI3K/Akt and Nrf2 pathways. *Int J Cardiol* 168: 1349-1359, 2013.
- Yang C, Zhang X, Fan H and Liu Y: Curcumin upregulates transcription factor Nrf2, HO-1 expression and protects rat brains against focal ischemia. *Brain Res* 1282: 133-141, 2009.